


# The Algebraic Counterpart of the Wagner Hierarchy

Jérémie Cabessa

University of Lausanne

July 3, 2008



## 1 Introduction

## 2 The Wagner hierarchy

## 3 $\omega$ -Semigroups

## 4 The FSG-hierarchy

## 5 Conclusion

# The Wagner Hierarchy

(most refined classification of  $\omega$ -rational languages)

Given two  $\omega$ -rational languages  $K$  and  $L$ , we set

$K \leq_w L$  iff  $K = f^{-1}(L)$  for some continuous function  $f$   
 iff Player II has a w.s. in the Wadge game  $W(K, L)$

Then as usual,  $K <_w L$  iff  $K \leq_w L$  and  $K \not\leq_w L$  and  $K \equiv_w L$  iff  $K \leq_w L \leq_w K$ .

## Definition (Wagner hierarchy)

The collection of all  $\omega$ -rational languages ordered by the relation  $\leq_w$  is called *the Wagner hierarchy*.

## The Wagner Hierarchy

(most refined classification of  $\omega$ -rational languages)

Given two  $\omega$ -rational languages  $K$  and  $L$ , we set

$$K \leq_w L \quad \text{iff} \quad K = f^{-1}(L) \quad \text{for some continuous function } f$$

$$\text{iff} \quad \text{Player II has a w.s. in the Wadge game } W(K, L)$$

Then as usual,  $K <_w L$  iff  $K \leq_w L$  and  $K \not\equiv_w L$  and  $K \equiv_w L$  iff  $K \leq_w L \leq_w K$ .

### Definition (Wagner hierarchy)

The collection of all  $\omega$ -rational languages ordered by the relation  $\leq_w$  is called *the Wagner hierarchy*.

## The Wagner Hierarchy

(most refined classification of  $\omega$ -rational languages)

Given two  $\omega$ -rational languages  $K$  and  $L$ , we set

$K \leq_w L$  iff  $K = f^{-1}(L)$  for some continuous function  $f$   
 iff Player II has a w.s. in the Wadge game  $W(K, L)$

Then as usual,  $K <_w L$  iff  $K \leq_w L$  and  $K \not\equiv_w L$  and  $K \equiv_w L$  iff  $K \leq_w L \leq_w K$ .

### Definition (Wagner hierarchy)

The collection of all  $\omega$ -rational languages ordered by the relation  $\leq_w$  is called *the Wagner hierarchy*.

## The Wagner Hierarchy

(most refined classification of  $\omega$ -rational languages)

Given two  $\omega$ -rational languages  $K$  and  $L$ , we set

$K \leq_w L$  iff  $K = f^{-1}(L)$  for some continuous function  $f$   
 iff Player II has a w.s. in the Wadge game  $W(K, L)$

Then as usual,  $K <_w L$  iff  $K \leq_w L \not\leq_w K$  and  $K \equiv_w L$  iff  $K \leq_w L \leq_w K$ .

### Definition (Wagner hierarchy)

The collection of all  $\omega$ -rational languages ordered by the relation  $\leq_w$  is called *the Wagner hierarchy*.

## The Wagner Hierarchy

(most refined classification of  $\omega$ -rational languages)

Given two  $\omega$ -rational languages  $K$  and  $L$ , we set

$K \leq_w L$  iff  $K = f^{-1}(L)$  for some continuous function  $f$   
 iff Player II has a w.s. in the Wadge game  $W(K, L)$

Then as usual,  $K <_w L$  iff  $K \leq_w L$  and  $K \not\equiv_w L$  and  $K \equiv_w L$  iff  $K \leq_w L \leq_w K$ .

### Definition (Wagner hierarchy)

The collection of all  $\omega$ -rational languages ordered by the relation  $\leq_w$  is called *the Wagner hierarchy*.




## Theorem (Wagner 79)

*The Wagner hierarchy has height  $\omega^\omega$  and it is decidable.*




## Theorem (Wagner 79)

*The Wagner hierarchy has height  $\omega^\omega$  and it is decidable.*




Let  $L$  be recognized by the following Muller automata




$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Let  $L$  be recognized by the following Muller automata




$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Let  $L$  be recognized by the following Muller automata




$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Let  $L$  be recognized by the following Muller automata



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$


Let  $L$  be recognized by the following Muller automata



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$




Let  $L$  be recognized by the following Muller automata



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Let  $L$  be recognized by the following Muller automata



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

## Theorem (Wagner)

*Let  $L$  be an  $\omega$ -rational language and  $\alpha < \omega^\omega$ . Then  $d_W(L) = \alpha$  iff any Muller  $\mathcal{A}(L)$  automata recognizing  $L$  contains a maximal  $\alpha$ -alternating tree.*


## 1 Introduction


## 2 The Wagner hierarchy

3  $\omega$ -Semigroups

## 4 The FSG-hierarchy

## 5 Conclusion





## Definition ( $\omega$ -semigroup)

An  $\omega$ -semigroup  $S$  is a pair  $(S_+, S_\omega)$ , where  $S_+$  is a semigroup,  $S_\omega$  is a set, and equipped with

- an associative mixed product :  $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product  $\pi_S : S_+^\omega \longrightarrow S_\omega$  satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

A *pointed  $\omega$ -semigroup* is a pair  $(S, X)$ , where

- $S = (S_+, S_\omega)$  is an  $\omega$ -semigroup,
- $X \subseteq S_\omega$ .

## Definition ( $\omega$ -semigroup)

An  $\omega$ -semigroup  $S$  is a pair  $(S_+, S_\omega)$ , where  $S_+$  is a semigroup,  $S_\omega$  is a set, and equipped with

- an associative mixed product :  $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product  $\pi_S : S_+^\omega \longrightarrow S_\omega$  satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

A *pointed  $\omega$ -semigroup* is a pair  $(S, X)$ , where

- $S = (S_+, S_\omega)$  is an  $\omega$ -semigroup,
- $X \subseteq S_\omega$ .



## Definition ( $\omega$ -semigroup)

An  $\omega$ -semigroup  $S$  is a pair  $(S_+, S_\omega)$ , where  $S_+$  is a semigroup,  $S_\omega$  is a set, and equipped with

- an associative mixed product :  $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product  $\pi_S : S_+^\omega \longrightarrow S_\omega$  satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

A *pointed  $\omega$ -semigroup* is a pair  $(S, X)$ , where

- $S = (S_+, S_\omega)$  is an  $\omega$ -semigroup,
- $X \subseteq S_\omega$ .

## Definition ( $\omega$ -semigroup)

An  $\omega$ -semigroup  $S$  is a pair  $(S_+, S_\omega)$ , where  $S_+$  is a semigroup,  $S_\omega$  is a set, and equipped with

- an associative mixed product :  $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product  $\pi_S : S_+^\omega \longrightarrow S_\omega$  satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

A *pointed  $\omega$ -semigroup* is a pair  $(S, X)$ , where

- $S = (S_+, S_\omega)$  is an  $\omega$ -semigroup,
- $X \subseteq S_\omega$ .

## Example

Let  $A$  be an alphabet. Then  $A^\infty = (A^+, A^\omega)$  equipped with the usual concatenation is an  $\omega$ -semigroup, called the *free  $\omega$ -semigroup* over  $A$ .

## Example

Let  $A$  be an alphabet. Then  $A^\infty = (A^+, A^\omega)$  equipped with the usual concatenation is an  $\omega$ -semigroup, called the *free  $\omega$ -semigroup* over  $A$ .

## Recognizability

$$(A^\infty, \textcolor{red}{L} \subseteq A^\omega)$$

## Example

Let  $A$  be an alphabet. Then  $A^\infty = (A^+, A^\omega)$  equipped with the usual concatenation is an  $\omega$ -semigroup, called the *free  $\omega$ -semigroup* over  $A$ .


## Recognizability

$$(A^\infty, \textcolor{red}{L} \subseteq A^\omega) \xrightarrow{\varphi} (S, X)$$

## Example

Let  $A$  be an alphabet. Then  $A^\infty = (A^+, A^\omega)$  equipped with the usual concatenation is an  $\omega$ -semigroup, called the *free  $\omega$ -semigroup* over  $A$ .

## Recognizability



## Finite pointed $\omega$ -semigroups are the algebraic counterparts of Büchi automata.

### Theorem

*An  $\omega$ -language is recognizable by a finite pointed  $\omega$ -semigroup iff it is recognizable by a finite Büchi (or Muller) automaton (iff it is  $\omega$ -rational).*


## Finite pointed $\omega$ -semigroups are the algebraic counterparts of Büchi automata.

### Theorem

*An  $\omega$ -language is recognizable by a finite pointed  $\omega$ -semigroup iff it is recognizable by a finite Büchi (or Muller) automaton (iff it is  $\omega$ -rational).*



Among all finite pointed  $\omega$ -semigroups recognizing a given  $\omega$ -rational language  $L$ , there exists a minimal one, **the syntactic  $\omega$ -semigroup** of  $L$ , denoted by  $Synt(L)$ .



## Example

Consider the language  $K = ((a + b)^*a)^\omega$ . Then

$\text{Synt}(K) = (S, X)$ , where

■  $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$  defined by the relations

$$0 \cdot 0 = 0$$

$$0 \cdot 1 = 0$$

$$1 \cdot 0 = 0$$

$$1 \cdot 1 = 1$$

$$00^\omega = 0^\omega$$

$$10^\omega = 0^\omega$$

$$01^\omega = 1^\omega$$

$$11^\omega = 1^\omega$$

■  $X = \{0^\omega\}$ .

## Example

Consider the language  $K = ((a + b)^*a)^\omega$ . Then  $\text{Synt}(K) = (S, X)$ , where

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$  defined by the relations

$$\begin{array}{cccc} 0 \cdot 0 = 0 & 0 \cdot 1 = 0 & 1 \cdot 0 = 0 & 1 \cdot 1 = 1 \\ 00^\omega = 0^\omega & 10^\omega = 0^\omega & 01^\omega = 1^\omega & 11^\omega = 1^\omega \end{array}$$

- $X = \{0^\omega\}$ .

## 1 Introduction

## 2 The Wagner hierarchy

## 3 $\omega$ -Semigroups

## 4 The FSG-hierarchy

## 5 Conclusion

We aim to classify finite pointed  $\omega$ -semigroups. We adopt a hierarchical game approach.

We transpose Wadge games on  $\omega$ -semigroups...

We aim to classify finite pointed  $\omega$ -semigroups. We adopt a hierarchical game approach.

**We transpose Wadge games on  $\omega$ -semigroups...**

Let  $S = (S_+, S_\omega)$  and  $T = (T_+, T_\omega)$  be two finite  $\omega$ -semigroups, and let also  $X \subseteq S_\omega$  and  $Y \subseteq T_\omega$ .

The infinite two-player game  $\text{SG}((S; X), (T, Y))$  is defined as follows:

- Player I plays elements from  $S_+$ ,
- Player II plays elements from  $T_+$ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let  $S = (S_+, S_\omega)$  and  $T = (T_+, T_\omega)$  be two finite  $\omega$ -semigroups, and let also  $X \subseteq S_\omega$  and  $Y \subseteq T_\omega$ .

The infinite two-player game  $\text{SG}((S; X), (T, Y))$  is defined as follows:

- Player I plays elements from  $S_+$ ,
- Player II plays elements from  $T_+$ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,



Let  $S = (S_+, S_\omega)$  and  $T = (T_+, T_\omega)$  be two finite  $\omega$ -semigroups, and let also  $X \subseteq S_\omega$  and  $Y \subseteq T_\omega$ .

The infinite two-player game  $\text{SG}((S; X), (T, Y))$  is defined as follows:

- Player I plays elements from  $S_+$ ,
- Player II plays elements from  $T_+$ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let  $S = (S_+, S_\omega)$  and  $T = (T_+, T_\omega)$  be two finite  $\omega$ -semigroups, and let also  $X \subseteq S_\omega$  and  $Y \subseteq T_\omega$ .

The infinite two-player game  $\text{SG}((S; X), (T; Y))$  is defined as follows:

- Player I plays elements from  $S_+$ ,
- Player II plays elements from  $T_+$ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let  $S = (S_+, S_\omega)$  and  $T = (T_+, T_\omega)$  be two finite  $\omega$ -semigroups, and let also  $X \subseteq S_\omega$  and  $Y \subseteq T_\omega$ .

The infinite two-player game  $\text{SG}((S; X), (T; Y))$  is defined as follows:

- Player I plays elements from  $S_+$ ,
- Player II plays elements from  $T_+$ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,



■ Player II wins the game  $\text{SG}(X, Y)$  iff

either  $\pi_S(s_0, s_1, \dots) \in X$  and  $\pi_T(t_0, t_1, \dots) \in Y$   
 or  $\pi_S(s_0, s_1, \dots) \notin X$  and  $\pi_T(t_0, t_1, \dots) \notin Y$

## Definition (SG-reduction)

We write  $(S, X) \leq_{\text{SG}} (T, Y)$  iff Player II has a winning strategy in  $\text{SG}(X, Y)$ . And as usual  $(S, X) \equiv_{\text{SG}} (T, Y)$  iff  $(S, X) \leq_{\text{SG}} (T, Y) \leq_{\text{SG}} (S, X)$ .



■ Player II wins the game  $\text{SG}(X, Y)$  iff

either  $\pi_S(s_0, s_1, \dots) \in X$  and  $\pi_T(t_0, t_1, \dots) \in Y$   
 or  $\pi_S(s_0, s_1, \dots) \notin X$  and  $\pi_T(t_0, t_1, \dots) \notin Y$

## Definition (SG-reduction)

We write  $(S, X) \leq_{\text{SG}} (T, Y)$  iff Player II has a winning strategy in  $\text{SG}(X, Y)$ . And as usual  $(S, X) \equiv_{\text{SG}} (T, Y)$  iff  $(S, X) \leq_{\text{SG}} (T, Y) \leq_{\text{SG}} (S, X)$ .



■ Player II wins the game  $\text{SG}(X, Y)$  iff

either  $\pi_S(s_0, s_1, \dots) \in X$  and  $\pi_T(t_0, t_1, \dots) \in Y$   
 or  $\pi_S(s_0, s_1, \dots) \notin X$  and  $\pi_T(t_0, t_1, \dots) \notin Y$

## Definition (SG-reduction)

We write  $(S, X) \leq_{\text{SG}} (T, Y)$  iff Player II has a winning strategy in  $\text{SG}(X, Y)$ . And as usual  $(S, X) \equiv_{\text{SG}} (T, Y)$  iff  $(S, X) \leq_{\text{SG}} (T, Y) \leq_{\text{SG}} (S, X)$ .



■ Player II wins the game  $\text{SG}(X, Y)$  iff

either  $\pi_S(s_0, s_1, \dots) \in X$  and  $\pi_T(t_0, t_1, \dots) \in Y$   
 or  $\pi_S(s_0, s_1, \dots) \notin X$  and  $\pi_T(t_0, t_1, \dots) \notin Y$

## Definition (SG-reduction)

We write  $(S, X) \leq_{\text{SG}} (T, Y)$  iff Player II has a winning strategy in  $\text{SG}(X, Y)$ . And as usual  $(S, X) \equiv_{\text{SG}} (T, Y)$  iff  $(S, X) \leq_{\text{SG}} (T, Y) \leq_{\text{SG}} (S, X)$ .

## Definition (FSG-hierarchy)

The collection of finite pointed  $\omega$ -semigroups ordered by the  $\leq_{SG}$ -relation is called *the FSG-hierarchy*.

## Theorem

*The FSG-hierarchy and the Wagner hierarchy are isomorphic.*






## Corollary

*The FSG-hierarchy has height  $\omega^\omega$ , and it is decidable.*

decidability procedure






*The FSG-hierarchy has height  $\omega^\omega$ , and it is decidable.*

The diagram illustrates the difference between the Wagner hierarchy and the FSG-hierarchy. On the left, the Wagner hierarchy is shown as a diamond-shaped graph with yellow nodes and solid black arrows. On the right, the FSG-hierarchy is shown as a similar diamond-shaped graph with blue nodes and solid black arrows. A red dashed arrow and three red question marks point to a missing edge between the top two nodes, indicating a difference in the structure. Below the FSG-hierarchy is the text  $Synt[(S,X)] \equiv_{sg} (S,X)$ .

# Corollary

*The FSG-hierarchy has height  $\omega^\omega$ , and it is decidable.*

decidability procedure



*The FSG-hierarchy has height  $\omega^\omega$ , and it is decidable.*

height  $\omega^0$

height  $\omega^0$

$Synt[(S,X)] \equiv_{sg} (S,X)$

The Wagner hierarchy

The FSG-hierarchy








## Corollary

*The FSG-hierarchy has height  $\omega^\omega$ , and it is decidable.*



## Corollary

*Let  $L$  be an  $\omega$ -rational language. Then  $d_W(L) = \alpha$  iff  $d_{FSG}(\text{Synt}(L)) = \alpha$ .*

## Example

Consider the syntactic pointed  $\omega$ -semigroup of  $K = ((a + b)^*a)^\omega$ :

■  $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$  defined by the relations

$$\begin{array}{cccc} 0 \cdot 0 = 0 & 0 \cdot 1 = 0 & 1 \cdot 0 = 0 & 1 \cdot 1 = 1 \\ 00^\omega = 0^\omega & 10^\omega = 0^\omega & 01^\omega = 1^\omega & 11^\omega = 1^\omega \end{array}$$

■  $X = \{0^\omega\}$ .

## Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$

$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$

●

1

●

0

Therefore  $d_{SG}((S, X)) = d_W(K) = \omega$ , and  $K$  is non-self-dual.

## Example (continued)

■  $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

■  $X = \{0^\omega\}.$




Therefore  $d_{SG}((S, X)) = d_W(K) = \omega$ , and  $K$  is non-self-dual.



## Example (continued)


- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ 
  - $0 \cdot 0 = 0$      $0 \cdot 1 = 0$
  - $1 \cdot 0 = 0$      $1 \cdot 1 = 1$
  - $00^\omega = 0^\omega$      $10^\omega = 0^\omega$
  - $01^\omega = 1^\omega$      $11^\omega = 1^\omega$
- $X = \{0^\omega\}$ .



## Example (continued)

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ 

$0 \cdot 0 = 0$	$0 \cdot 1 = 0$
$1 \cdot 0 = 0$	$1 \cdot 1 = 1$
$00^\omega = 0^\omega$	$10^\omega = 0^\omega$
$01^\omega = 1^\omega$	$11^\omega = 1^\omega$
- $X = \{0^\omega\}$ .



## Example (continued)

■  $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$


$0 \cdot 0 = 0$      $0 \cdot 1 = 0$

$1 \cdot 0 = 0$      $1 \cdot 1 = 1$

$00^\omega = 0^\omega$      $10^\omega = 0^\omega$

$01^\omega = 1^\omega$      $11^\omega = 1^\omega$

■  $X = \{0^\omega\}$ .




Therefore  $d_{SG}((S, X)) = d_W(K) = \omega$ , and  $K$  is non-self-dual.



### Example (continued)

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ 

$0 \cdot 0 = 0$	$0 \cdot 1 = 0$
$1 \cdot 0 = 0$	$1 \cdot 1 = 1$
$00^\omega = 0^\omega$	$10^\omega = 0^\omega$
$01^\omega = 1^\omega$	$11^\omega = 1^\omega$
- $X = \{0^\omega\}$ .



### Example (continued)

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

- $X = \{0^\omega\}$ .



## Example (continued)

■  $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

■  $X = \{0^\omega\}$ .



Therefore  $d_{SG}((S, X)) = d_W(K) = \omega$ , and  $K$  is non-self-dual.

## Example

Consider the syntactic pointed  $\omega$ -semigroup of

$$L = (a\{b, c\}^* \cup \{b\})^\omega:$$

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$  is defined by

$$a^2 = a$$

$$ab = a$$

$$ac = a$$

$$ba = a$$

$$b^2 = b$$

$$bc = c$$

$$cb = c$$

$$c^2 = c$$

$$b^\omega = a^\omega$$

$$c^\omega = 0$$

$$aa^\omega = a^\omega$$

$$a(ca)^\omega = a^\omega$$

$$ba^\omega = a^\omega$$

$$b(ca)^\omega = (ca)^\omega$$

$$ca^\omega = (ca)^\omega$$

$$c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$

## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

$$\blacksquare T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

$$\blacksquare Y = \{a^\omega\}.$$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$





## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



## Example (continued)

■  $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■  $Y = \{a^\omega\}.$



Therefore  $d_{SG}((T, Y)) = d_W(L) = \omega^2$ ,  
and  $L$  is non-self-dual.

## Proposition

*Let  $(S, X)$  be a finite pointed  $\omega$ -semigroup.  $d_{FSG}((S, X)) = \alpha$  iff  $(S, X)$  contains a maximal  $\alpha$ -alternating tree.*

## Corollary

*Let  $L$  be an  $\omega$ -rational language. Then  $d_W(L) = \alpha$  iff any finite pointed  $\omega$  semigroup recognizing  $L$  contains a maximal  $\alpha$ -alternating tree.*



## Proposition

*Let  $(S, X)$  be a finite pointed  $\omega$ -semigroup.  $d_{FSG}((S, X)) = \alpha$  iff  $(S, X)$  contains a maximal  $\alpha$ -alternating tree.*

## Corollary

*Let  $L$  be an  $\omega$ -rational language. Then  $d_W(L) = \alpha$  iff any finite pointed  $\omega$  semigroup recognizing  $L$  contains a maximal  $\alpha$ -alternating tree.*

## Summary :

### The Wagner hierarchy

From the **automata** point of view

#### Theorem

*Then  $d_W(L) = \alpha$  iff any Muller automata recognizing  $L$  contains a maximal  $\alpha$ -alternating tree.*

From the **algebraic** point of view

#### Theorem

*Then  $d_W(L) = \alpha$  iff any finite  $\omega$ -semigroup recognizing  $L$  contains a maximal  $\alpha$ -alternating tree.*

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- One can compute the Wagner degree of an  $\omega$ -rational language directly on its syntactic image.
- Build a finite pointed  $\omega$ -semigroup of any given Wagner degree.

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- One can compute the Wagner degree of an  $\omega$ -rational language directly on its syntactic image.
- Build a finite pointed  $\omega$ -semigroup of any given Wagner degree.

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- **One can compute the Wagner degree of an  $\omega$ -rational language directly on its syntactic image.**
- Build a finite pointed  $\omega$ -semigroup of any given Wagner degree.

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- **One can compute the Wagner degree of an  $\omega$ -rational language directly on its syntactic image.**
- Build a finite pointed  $\omega$ -semigroup of any given Wagner degree.