

Introduction to Topology

Jérémie Cabessa

UNIL Institute bdbdbdb

11th July 2007

Outline

- ① Metric spaces
- ② Topological spaces
- ③ Compactness
- ④ Connectedness

Metric spaces

A *metric space* is a set of points and a distance which measures of the degree of closeness of pairs of points in this space.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Definition

A metric space is a pair (X, d) , where X is a set and d is a distance on X , i.e. d is a function $d : X \times X \longrightarrow \mathbb{R}$ such that for all $x, y, z \in X$,

- ① $d(x, y) \geq 0$,
- ② $d(x, y) = 0$ if and only if $x = y$,
- ③ $d(x, y) = d(y, x)$,
- ④ $d(x, z) \leq d(x, y) + d(y, z)$.

Example

- ① (\mathbb{R}, d) is a metric space, where d is defined by $d(a, b) = |b - a|$.
- ② (\mathbb{R}^n, d_n) is a metric space, where d_n is defined by $d(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$.

Example


- ① (\mathbb{R}, d) is a metric space, where d is defined by $d(a, b) = |b - a|$.
- ② (\mathbb{R}^n, d_n) is a metric space, where d_n is defined by $d(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$.

Example

- ① (\mathbb{R}, d) is a metric space, where d is defined by $d(a, b) = |b - a|$.
- ② (\mathbb{R}^n, d_n) is a metric space, where d_n is defined by $d(x, y) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$.

Example


For $n = 2$, this distance is the classical distance used in analytical geometry:




Example

Let A be an alphabet. Then (A^ω, d) is a metric space, with d is defined by $d(\alpha, \beta) = 2^{-r}$, where $r = \min\{n : \alpha(n) \neq \beta(n)\}$.

A function f from a metric space X to another metric space Y is *continuous at the point a* if for any (small) “neighborhood” of $f(a)$, denoted by V , there is a “neighborhood” of a , denoted by U , such that if $f(U) \subseteq V$.



The following function is not continuous at the point a .



By choosing such a V , there is no U such that $f(U) \subset V$.

Definition

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in X$, if $d(x, a) < \delta$ then $d'(f(x), f(a)) < \varepsilon$.

A function $f : X \longrightarrow Y$ is continuous iff it is continuous at each point of X .

Definition

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in X$, if $d(x, a) < \delta$ then $d'(f(x), f(a)) < \varepsilon$.

A function $f : X \longrightarrow Y$ is continuous iff it is continuous at each point of X .

Definition

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in X$, if $d(x, a) < \delta$ then $d'(f(x), f(a)) < \varepsilon$.

A function $f : X \longrightarrow Y$ is continuous iff it is continuous at each point of X .

Definition

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x \in X$, if $d(x, a) < \delta$ then $d'(f(x), f(a)) < \varepsilon$.

A function $f : X \longrightarrow Y$ is continuous iff it is continuous at each point of X .

A function $f : X \longrightarrow Y$ is continuous at the point a iff

$$\forall \varepsilon \exists \delta \forall x (P \Rightarrow Q).$$

A function $f : X \longrightarrow Y$ is not continuous at the point a iff

$$\exists \varepsilon \forall \delta \exists x (P \wedge \neg Q).$$

A function $f : X \longrightarrow Y$ is continuous at the point a iff

$$\forall \varepsilon \exists \delta \forall x (P \Rightarrow Q).$$

A function $f : X \longrightarrow Y$ is not continuous at the point a iff

$$\exists \varepsilon \forall \delta \exists x (P \wedge \neg Q).$$

A function $f : X \longrightarrow Y$ is continuous at the point a iff

$$\forall \varepsilon \exists \delta \forall x (P \Rightarrow Q).$$

A function $f : X \longrightarrow Y$ is not continuous at the point a iff

$$\exists \varepsilon \forall \delta \exists x (P \wedge \neg Q).$$

A function $f : X \longrightarrow Y$ is continuous at the point a iff

$$\forall \varepsilon \exists \delta \forall x (P \Rightarrow Q).$$

A function $f : X \longrightarrow Y$ is not continuous at the point a iff

$$\exists \varepsilon \forall \delta \exists x (P \wedge \neg Q).$$

Example

- 1 Let $f : (X, d) \longrightarrow (X', d')$ be a constant function, then f is continuous. Take any δ , say $\delta = 1$.
- 2 The identity function $id : (X, d) \longrightarrow (X, d)$ is continuous. Take $\delta = \varepsilon$.


Example

- 1 Let $f : (X, d) \longrightarrow (X', d')$ be a constant function, then f is continuous. Take any δ , say $\delta = 1$.
- 2 The identity function $id : (X, d) \longrightarrow (X, d)$ is continuous. Take $\delta = \varepsilon$.

Definition

Let (X, d) be a metric space, and let $a \in X$ and δ be given. The open sphere about a of radius δ is the set

$$S(a; \delta) = \{x \in X : d(a, x) < \delta\}.$$



One has by definition

$$x \in S(a; \delta) \text{ iff } d(a, x) < \delta.$$

Therefore the definition of the continuity becomes...

Theorem

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that

$$f(S(a; \delta)) \subset S(f(a); \varepsilon).$$

Therefore the definition of the continuity becomes...

Theorem

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that

$$f(S(a; \delta)) \subset S(f(a); \varepsilon).$$

And the property $f(A) \subset B$ iff $A \subset f^{-1}(B)$ implies...

Theorem

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that

$$S(a; \delta) \subset f^{-1}(S(f(a); \varepsilon)).$$

And the property $f(A) \subset B$ iff $A \subset f^{-1}(B)$
implies...

Theorem


Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for all $\varepsilon > 0$ there exists $\delta > 0$ such that

$$S(a; \delta) \subset f^{-1}(S(f(a); \varepsilon)).$$

Definition

Let (X, d) and let $a \in X$. A subset N of X is called a neighborhood of a if there exists a $\delta > 0$ such that

$$S(a; \delta) \subset N.$$



Theorem

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for every neighborhood N of $f(a)$ there exists a corresponding neighborhood M of a such that

$$f(M) \subset N,$$

or equivalently

$$M \subset f^{-1}(N).$$

Theorem

Let (X, d) and (X', d') be two metric spaces, and let $a \in X$. A function $f : X \longrightarrow Y$ is continuous at the point a iff for every neighborhood N of $f(a)$ there exists a corresponding neighborhood M of a such that

$$f(M) \subset N,$$

or equivalently

$$M \subset f^{-1}(N).$$


Proof.

On the blackboard...



Definition

A subset O of a metric space is said to be open if it is a neighborhood of each of its point.



Example

- 1 Every “open interval” $]a, b[$ is an open set of \mathbb{R} (equipped with the usual metric).

$$]a, b[= \{x \in \mathbb{R} : a < x < b\},$$

- 2 Every cross product of “open intervals” $]a_i, b_i[$ is an open set of \mathbb{R}^n (equipped with the usual metric).

$$\prod_{i=1}^n]a_i, b_i[.$$

Example

- 1 Every “open interval” $]a, b[$ is an open set of \mathbb{R} (equipped with the usual metric).

$$]a, b[= \{x \in \mathbb{R} : a < x < b\},$$

- 2 Every cross product of “open intervals” $]a_i, b_i[$ is an open set of \mathbb{R}^n (equipped with the usual metric).


$$\prod_{i=1}^n]a_i, b_i[.$$

Lemma

Every open sphere is open.

Proof.


Let $S(a; \delta)$ be an open sphere. We need to prove that $S(a; \delta)$ is a neighborhood of each of its point. Let $b \in S(a; \delta)$ and let $\varepsilon < \delta - d(a, b)$. Then $S(b; \varepsilon) \subseteq S(a; \delta)$, which concludes the proof.



Proof.


Let $S(a; \delta)$ be an open sphere. We need to prove that $S(a; \delta)$ is a neighborhood of each of its point.

Let $b \in S(a; \delta)$ and let $\varepsilon < \delta - d(a, b)$. Then $S(b; \varepsilon) \subseteq S(a; \delta)$, which concludes the proof.




Proof.

Let $S(a; \delta)$ be an open sphere. We need to prove that $S(a; \delta)$ is a neighborhood of each of its point. Let $b \in S(a; \delta)$ and let $\varepsilon < \delta - d(a, b)$. Then $S(b; \varepsilon) \subseteq S(a; \delta)$, which concludes the proof.



Proof.

Let $S(a; \delta)$ be an open sphere. We need to prove that $S(a; \delta)$ is a neighborhood of each of its point. Let $b \in S(a; \delta)$ and let $\varepsilon < \delta - d(a, b)$. Then $S(b; \varepsilon) \subseteq S(a; \delta)$, which concludes the proof.



Theorem

Let (X, d) and (X', d') be two metric spaces and let $f : (X, d) \longrightarrow (X', d')$. Then f is continuous if and only if the inverse image of every open set of X' is an open set of X .

Proof.

On the blackboard. □

Topological spaces

Definition

A *topological space* is a pair (X, \mathcal{J}) , where X is a non-empty set and \mathcal{J} is a collection of subsets of X such that

- 1 $X, \emptyset \in \mathcal{J}$,
- 2 If $O_1, \dots, O_n \in \mathcal{J}$, then $O_1 \cap O_2 \cap \dots \cap O_n \in \mathcal{J}$,
- 3 If $O_\alpha \in \mathcal{J}$ for each $\alpha \in I$, then $\bigcup_{\alpha \in I} O_\alpha \in \mathcal{J}$.

Definition

A *topological space* is a pair (X, \mathcal{J}) , where X is a non-empty set and \mathcal{J} is a collection of subsets of X such that

- 1 $X, \emptyset \in \mathcal{J}$,
- 2 If $O_1, \dots, O_n \in \mathcal{J}$, then
 $O_1 \cap O_2 \cap \dots \cap O_n \in \mathcal{J}$,
- 3 If $O_\alpha \in \mathcal{J}$ for each $\alpha \in I$, then $\bigcup_{\alpha \in I} O_\alpha \in \mathcal{J}$.

Definition

A *topological space* is a pair (X, \mathcal{J}) , where X is a non-empty set and \mathcal{J} is a collection of subsets of X such that

- ① $X, \emptyset \in \mathcal{J}$,
- ② If $O_1, \dots, O_n \in \mathcal{J}$, then
 $O_1 \cap O_2 \cap \dots \cap O_n \in \mathcal{J}$,
- ③ If $O_\alpha \in \mathcal{J}$ for each $\alpha \in I$, then $\bigcup_{\alpha \in I} O_\alpha \in \mathcal{J}$.

Definition

A *topological space* is a pair (X, \mathcal{J}) , where X is a non-empty set and \mathcal{J} is a collection of subsets of X such that

- ① $X, \emptyset \in \mathcal{J}$,
- ② If $O_1, \dots, O_n \in \mathcal{J}$, then
 $O_1 \cap O_2 \cap \dots \cap O_n \in \mathcal{J}$,
- ③ If $O_\alpha \in \mathcal{J}$ for each $\alpha \in I$, then $\bigcup_{\alpha \in I} O_\alpha \in \mathcal{J}$.

Elements of the collection \mathcal{J} are called the *open sets* of X .

A subset is said to be *closed* if it is the complement of an open set.

Elements of the collection \mathcal{J} are called the *open sets* of X .

A subset is said to be *closed* if it is the complement of an open set.

Example

Let (X, \mathcal{T}) be a topological space.

- \emptyset is closed, since $\emptyset^c = X$ is open.
- X is closed, since $X^c = \emptyset$ is open.
- Therefore \emptyset and X are *clopen* sets.

Example

Let (X, \mathcal{T}) be a topological space.

- \emptyset is closed, since $\emptyset^c = X$ is open.
- X is closed, since $X^c = \emptyset$ is open.
- Therefore \emptyset and X are *clopen* sets.

Example

Let (X, \mathcal{T}) be a topological space.

- \emptyset is closed, since $\emptyset^c = X$ is open.
- X is closed, since $X^c = \emptyset$ is open.
- Therefore \emptyset and X are *clopen* sets.

Example

Let (X, \mathcal{T}) be a topological space.

- \emptyset is closed, since $\emptyset^c = X$ is open.
- X is closed, since $X^c = \emptyset$ is open.
- Therefore \emptyset and X are *clopen* sets.

From metric spaces to topological spaces...

Let (X, d) be a metric space and let \mathcal{J} be the collection of open sets of X , Then (X, \mathcal{J}) is a topological space.

From metric spaces to topological spaces...

Let (X, d) be a metric space and let \mathcal{J} be the collection of open sets of X , Then (X, \mathcal{J}) is a topological space.

Example

- 1 The *real line* is the topological space arising from the metric space (\mathbb{R}, d) , where $d(a, b) = |b - a|$.
- 2 The *Euclidian n -space with the usual topology* is the topological space arising from the metric space (\mathbb{R}^n, d) .

Example

- ① The *real line* is the topological space arising from the metric space (\mathbb{R}, d) , where $d(a, b) = |b - a|$.
- ② The *Euclidian n -space with the usual topology* is the topological space arising from the metric space (\mathbb{R}^n, d) .

Example

- 1 Let X be a non-empty arbitrary set. Let $\mathcal{J} = \{\emptyset, X\}$. Then (X, \mathcal{J}) is a topological space.
- 2 Let X be a non-empty arbitrary set. Then $(X, \mathcal{P}(X))$ is a topological space. This topology on X is the one which contains the largest number of elements. It is called the *discrete topology*.

Example

- ① Let X be a non-empty arbitrary set. Let $\mathcal{J} = \{\emptyset, X\}$. Then (X, \mathcal{J}) is a topological space.
- ② Let X be a non-empty arbitrary set. Then $(X, \mathcal{P}(X))$ is a topological space. This topology on X is the one which contains the largest number of elements. It is called the *discrete topology*.

Example

Let $X = \{a, b\}$. Let $\mathcal{J}_1 = \{\emptyset, X\}$,
 $\mathcal{J}_2 = \{\emptyset, \{a\}, X\}$, $\mathcal{J}_3 = \{\emptyset, \{b\}, X\}$,
 $\mathcal{J}_4 = \{\emptyset, \{a\}, \{b\}, X\}$. Then (X, \mathcal{J}_i) , for
 $i = 1, 2, 3, 4$, are four distinct topological spaces.

Example


Let \mathbb{N} be the set of positive integers. For each $n \in \mathbb{N}$, let $O_n = \{n, n+1, n+2, \dots\}$, and let $\mathcal{J} = \{\emptyset, O_0, O_1, O_2, \dots\}$. Then $(\mathbb{N}, \mathcal{J})$ is a topological space.

Definition

Let (X, \mathcal{J}) and (X', \mathcal{J}') be two topological spaces and let $f : X \longrightarrow Y$. Then f is said to be *continuous* if and only if the preimage of any open set is an open set.

Definition

The *closure* \bar{A} of a subset A is the smallest closed set containing A .



Lemma

A is closed if and only if $A = \bar{A}$.

Proof.

(\Rightarrow) Suppose that A is closed. Then the smallest closed set containing A is A itself, i.e $\bar{A} = A$.

(\Leftarrow) Suppose $A = \bar{A}$. Since \bar{A} is closed by definition, then A is also closed.



Lemma

A is closed if and only if $A = \bar{A}$.

Proof.

(\Rightarrow) Suppose that A is closed. Then the smallest closed set containing A is A itself, i.e $\bar{A} = A$.

(\Leftarrow) Suppose $A = \bar{A}$. Since \bar{A} is closed by definition, then A is also closed.



Lemma

A is closed if and only if $A = \bar{A}$.


Proof.

- (\Rightarrow) Suppose that A is closed. Then the smallest closed set containing A is A itself, i.e $\bar{A} = A$.
- (\Leftarrow) Suppose $A = \bar{A}$. Since \bar{A} is closed by definition, then A is also closed.



Definition


The *interior* $\text{int}(A)$ of a subset A is the largest open set contained in A .



Compactness


A *compact* set is a closed and bounded set.

Bounded in the sense that it can be covered with a finite number of open sets.



Compact set

A *compact* set is a closed and bounded set.
Bounded in the sense that it can be covered with
a finite number of open sets.



Compact set

Definition

Let X be a set and let $(A_i)_{i \in \alpha}$ be a family of sets. The collection $(A_i)_{i \in \alpha}$ is called a *covering* of X if $X \subseteq \bigcup_{i \in \alpha} A_i$.

Given $\beta \subseteq \alpha$, if the the subfamily $(A_i)_{i \in \beta}$ still covers X , then $(A_i)_{i \in \beta}$ is called a *subcovering* of $(A_i)_{i \in \alpha}$.

If A_i is open for each $i \in \alpha$, then $(A_i)_{i \in \alpha}$ is called an *open covering* of X .

Definition

Let X be a set and let $(A_i)_{i \in \alpha}$ be a family of sets. The collection $(A_i)_{i \in \alpha}$ is called a *covering* of X if $X \subseteq \bigcup_{i \in \alpha} A_i$.

Given $\beta \subseteq \alpha$, if the the subfamily $(A_i)_{i \in \beta}$ still covers X , then $(A_i)_{i \in \beta}$ is called a *subcovering* of $(A_i)_{i \in \alpha}$.

If A_i is open for each $i \in \alpha$, then $(A_i)_{i \in \alpha}$ is called an *open covering* of X .

Definition


Let X be a set and let $(A_i)_{i \in \alpha}$ be a family of sets. The collection $(A_i)_{i \in \alpha}$ is called a *covering* of X if $X \subseteq \bigcup_{i \in \alpha} A_i$.

Given $\beta \subseteq \alpha$, if the the subfamily $(A_i)_{i \in \beta}$ still covers X , then $(A_i)_{i \in \beta}$ is called a *subcovering* of $(A_i)_{i \in \alpha}$.

If A_i is open for each $i \in \alpha$, then $(A_i)_{i \in \alpha}$ is called an *open covering* of X .

Example

For each integer n , let $A_n = [n, n + 1]$. Then $(A_n)_{n \in \mathbb{Z}}$ is an infinite covering of \mathbb{R} .



Definition

A topological space (X, \mathcal{J}) is called *compact* iff from each open covering of X , one may extract a finite subcovering.

Definition

A subset A of (X, \mathcal{J}) is called *compact* iff from each open covering of A , one may extract a finite subcovering.

Definition

A topological space (X, \mathcal{J}) is called *compact* iff from each open covering of X , one may extract a finite subcovering.

Definition

A subset A of (X, \mathcal{J}) is called *compact* iff from each open covering of A , one may extract a finite subcovering.

Lemma

The closed interval $[0, 1]$ is compact.

Towards a contradiction, suppose that $[0, 1]$ is not compact. Then there exists an open covering $(O_\alpha)_{\alpha \in I}$ of $[0, 1]$ from which one cannot extract a finite subcovering of $[0, 1]$.


Towards a contradiction, suppose that $[0, 1]$ is not compact. Then there exists an open covering $(O_\alpha)_{\alpha \in I}$ of $[0, 1]$ from which one cannot extract a finite subcovering of $[0, 1]$.

Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:

Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:




Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:



Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:



Therefore, we can build the following infinite sequence of intervals $[a_n, b_n]$, for any $n > 0$:



And so on, and so forth.

For each $n > 0$, one has

- $|b_n - a_n| = \frac{1}{2^n},$
 - $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n.$
- (1)

Now let a be the smallest upper bound of the $(a_i)_{i>0}$, and let b be the greatest lower bound of the $(b_i)_{i>0}$



For each $n > 0$, one has

- $|b_n - a_n| = \frac{1}{2^n},$
 - $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n.$
- (1)

Now let a be the smallest upper bound of the $(a_i)_{i>0}$, and let b be the greatest lower bound of the $(b_i)_{i>0}$



Then (1) implies $a_n \leq a \leq b \leq b_n$, for all $n > 0$.

Hence $|b - a| \leq \frac{1}{2^n}$, for all $n > 0$. Therefore $a = b$.

Then (1) implies $a_n \leq a \leq b \leq b_n$, for all $n > 0$.

Hence $|b - a| \leq \frac{1}{2^n}$, for all $n > 0$. Therefore $a = b$.

Since the covering $(O_\alpha)_{\alpha \in I}$ covers $[0, 1]$ and $a = b \in [0, 1]$, there exists an open set O_β such that $a \in O_\beta$.

Since O_β is open, there is an $\varepsilon > 0$ such that $S(a; \varepsilon) \subseteq O_\beta$.

Now let N large enough such that $\frac{1}{2^N} < \varepsilon$. Then $b_N - a_N = \frac{1}{2^N} < \varepsilon$.

Since the covering $(O_\alpha)_{\alpha \in I}$ covers $[0, 1]$ and $a = b \in [0, 1]$, there exists an open set O_β such that $a \in O_\beta$.

Since O_β is open, there is an $\varepsilon > 0$ such that $S(a; \varepsilon) \subseteq O_\beta$.


Now let N large enough such that $\frac{1}{2^N} < \varepsilon$. Then $b_N - a_N = \frac{1}{2^N} < \varepsilon$.

Since the covering $(O_\alpha)_{\alpha \in I}$ covers $[0, 1]$ and $a = b \in [0, 1]$, there exists an open set O_β such that $a \in O_\beta$.

Since O_β is open, there is an $\varepsilon > 0$ such that $S(a; \varepsilon) \subseteq O_\beta$.


Now let N large enough such that $\frac{1}{2^N} < \varepsilon$. Then $b_N - a_N = \frac{1}{2^N} < \varepsilon$.

One has $b_N - a_N < \varepsilon$ and $a = b \in [a_N, b_N]$.
Therefore $[a_N, b_N] \subseteq S(a; \varepsilon) \subseteq O_\beta$.



In other terms, $[a_N, b_N]$ is covered with a finite subcovering of $(O_\alpha)_{\alpha \in I}$, namely O_β itself. A contradiction.

One has $b_N - a_N < \varepsilon$ and $a = b \in [a_N, b_N]$.
 Therefore $[a_N, b_N] \subseteq S(a; \varepsilon) \subseteq O_\beta$.




In other terms, $[a_N, b_N]$ is covered with a finite subcovering of $(O_\alpha)_{\alpha \in I}$, namely O_β itself. A contradiction.

Corollary


Each closed interval $[a, b]$ is compact.

Connectedness

A topological space is *connected* if it is all of one piece. That is if it is impossible to decompose it into two disjoint non-empty open sets.




Connected space




Non-connected space

A topological space is *connected* if it is all of one piece. That is if it is impossible to decompose it into two disjoint non-empty open sets.



Connected space



Non-connected space

Definition

A topological space X is said to be *connected* if the only two clopen sets of X are \emptyset and X itself.

Lemma

A topological space X is not connected if and only if there are two non-empty open sets P and Q such that

- $P \cup Q = X$,
- $P \cap Q = \emptyset$.

Proof.

(\Rightarrow) If X is not connected, there exists a clopen set C such that $C \neq \emptyset, X$.

Therefore C and C^c are open sets, and $C \cup C^c = X$ and $C \cap C^c = \emptyset$.

(\Leftarrow) If there are two non-empty open sets P and Q such that $P \cup Q = X$ and $P \cap Q = \emptyset$, then obviously $P = Q^c$ and $Q = P^c$. These properties imply P and Q clopen different from \emptyset and X .



Proof.

(\Rightarrow) If X is not connected, there exists a clopen set C such that $C \neq \emptyset, X$.
Therefore C and C^c are open sets, and $C \cup C^c = X$ and $C \cap C^c = \emptyset$.

(\Leftarrow) If there are two non-empty open sets P and Q such that $P \cup Q = X$ and $P \cap Q = \emptyset$, then obviously $P = Q^c$ and $Q = P^c$. These properties imply P and Q clopen different from \emptyset and X .



Proof.

- (\Rightarrow) If X is not connected, there exists a clopen set C such that $C \neq \emptyset, X$.
Therefore C and C^c are open sets, and $C \cup C^c = X$ and $C \cap C^c = \emptyset$.
- (\Leftarrow) If there are two non-empty open sets P and Q such that $P \cup Q = X$ and $P \cap Q = \emptyset$, then obviously $P = Q^c$ and $Q = P^c$. These properties imply P and Q clopen different from \emptyset and X .



Proof.

- (\Rightarrow) If X is not connected, there exists a clopen set C such that $C \neq \emptyset, X$.
Therefore C and C^c are open sets, and $C \cup C^c = X$ and $C \cap C^c = \emptyset$.
- (\Leftarrow) If there are two non-empty open sets P and Q such that $P \cup Q = X$ and $P \cap Q = \emptyset$, then obviously $P = Q^c$ and $Q = P^c$. These properties imply P and Q clopen different from \emptyset and X .



One can prove that connected subsets of the real line \mathbb{R} are exactly the intervals.