

The Algebraic Counterpart of the Wagner Hierarchy

Jérémie Cabessa

University of Lausanne

January 24, 2008

- 1 Introduction
- 2 ω -Rational Languages
- 3 The Wagner hierarchy
- 4 ω -Semigroups
- 5 The FSG-hierarchy
- 6 Conclusion


- 1 Introduction
- 2 ω -Rational Languages
- 3 The Wagner hierarchy
- 4 ω -Semigroups
- 5 The FSG-hierarchy
- 6 Conclusion

Outline

- 1 Introduction
- 2 ω -Rational Languages
- 3 The Wagner hierarchy
- 4 ω -Semigroups
- 5 The FSG-hierarchy
- 6 Conclusion


Outline

- 1 Introduction
- 2 ω -Rational Languages
- 3 The Wagner hierarchy
- 4 ω -Semigroups
- 5 The FSG-hierarchy
- 6 Conclusion



Muller automaton (deterministic)

\mathcal{A} is a labeled graph, with $\mathcal{T}_{\mathcal{A}} \subseteq \mathcal{P}^{\mathcal{Q}}$.




$$\mathcal{T}_{\mathcal{A}} = \{\{q_0\}, \{q_1\}\}$$

Then $L^{\omega}(\mathcal{A}) = A^*(0^{\omega} \cup 1^{\omega})$.

Muller automaton (deterministic)

\mathcal{A} is a labeled graph, with $\mathcal{T}_{\mathcal{A}} \subseteq \mathcal{P}^{\omega}$.



$$\mathcal{T}_{\mathcal{A}} = \{\{q_0\}, \{q_1\}\}$$

Then $L^{\omega}(\mathcal{A}) = A^*(0^{\omega} \cup 1^{\omega})$.

The Wagner Hierarchy

(classification of ω -rational languages)

Given two ω -rational languages K and L , we set

$$K \leq_W L \quad \text{iff} \quad K = f^{-1}(L) \text{ for some continuous function } f$$

$$\text{iff} \quad w \in K \Leftrightarrow f(w) \in L.$$

$$K <_W L \quad \text{iff} \quad K \leq_W L \not\leq_W K$$

$$K \equiv_W L \quad \text{iff} \quad K \leq_W L \leq_W K.$$

Definition (Wagner hierarchy)

The collection of all ω -rational languages ordered by the relation \leq_W is called *the Wagner hierarchy*.

The Wagner Hierarchy

(classification of ω -rational languages)

Given two ω -rational languages K and L , we set

$$K \leq_W L \quad \text{iff} \quad K = f^{-1}(L) \text{ for some continuous function } f$$

$$\text{iff} \quad w \in K \Leftrightarrow f(w) \in L.$$

$$K <_W L \quad \text{iff} \quad K \leq_W L \not\leq_W K$$

$$K \equiv_W L \quad \text{iff} \quad K \leq_W L \leq_W K.$$

Definition (Wagner hierarchy)

The collection of all ω -rational languages ordered by the relation \leq_W is called *the Wagner hierarchy*.

The Wagner Hierarchy

(classification of ω -rational languages)

Given two ω -rational languages K and L , we set

$$K \leq_W L \quad \text{iff} \quad K = f^{-1}(L) \text{ for some continuous function } f$$

$$\text{iff} \quad w \in K \Leftrightarrow f(w) \in L.$$

$$K <_W L \quad \text{iff} \quad K \leq_W L \not\leq_W K$$

$$K \equiv_W L \quad \text{iff} \quad K \leq_W L \leq_W K.$$

Definition (Wagner hierarchy)

The collection of all ω -rational languages ordered by the relation \leq_W is called *the Wagner hierarchy*.

The Wagner Hierarchy

(classification of ω -rational languages)

Given two ω -rational languages K and L , we set

$K \leq_W L$ iff $K = f^{-1}(L)$ for some continuous function f

iff $w \in K \Leftrightarrow f(w) \in L$.

$K <_W L$ iff $K \leq_W L \not\leq_W K$

$K \equiv_W L$ iff $K \leq_W L \leq_W K$.

Definition (Wagner hierarchy)

The collection of all ω -rational languages ordered by the relation \leq_W is called *the Wagner hierarchy*.

The Wagner Hierarchy

(classification of ω -rational languages)

Given two ω -rational languages K and L , we set

$$K \leq_W L \quad \text{iff} \quad K = f^{-1}(L) \text{ for some continuous function } f$$

$$\text{iff} \quad w \in K \Leftrightarrow f(w) \in L.$$

$$K <_W L \quad \text{iff} \quad K \leq_W L \not\leq_W K$$


$$K \equiv_W L \quad \text{iff} \quad K \leq_W L \leq_W K.$$

Definition (Wagner hierarchy)

The collection of all ω -rational languages ordered by the relation \leq_W is called *the Wagner hierarchy*.

Theorem

The Wagner hierarchy has height ω^ω and it is decidable.



Wagner's decidability procedure

Classifying ω -rational language is equivalent to classifying their underlying Muller automata.

/

$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}\}, \{q_{11}\}\}$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.

/

$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}\}, \{q_{11}\}\}$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\} \equiv \dots$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\}$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\} \equiv \dots$$

Wagner's decidability procedure


Classifying ω -rational language is equivalent to classifying their underlying Muller automata.




$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\} \equiv \dots$$


Wagner's decidability procedure

Classifying ω -rational language is equivalent to classifying their underlying Muller automata.



$$\mathcal{T} = \{\{q_0\}, \{q_2, q_3\}, \{q_4\}, \{q_6, q_7\}, \{q_8\}, \{q_{10}, q_{11}\}, \{q_{12}\}\} \equiv \dots$$





A semigroup (S, \cdot) is a set S equipped with an associative binary operation “ \cdot ” on S .

Example

The set $\{0, 1\}$ equipped with the usual multiplication is a semigroup.

A semigroup (S, \cdot) is a set S equipped with an associative binary operation “ \cdot ” on S .

Example

The set $\{0, 1\}$ equipped with the usual multiplication is a semigroup.

An ω -semigroup is a semigroup equipped with a suitable infinite product.

Definition (ω -semigroup)

An ω -semigroup S is a pair (S_+, S_ω) , where S_+ is a semigroup, S_ω is a set, and equipped with

- an associative mixed product : $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product $\pi_S : S_+^\omega \longrightarrow S_\omega$ satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

Notation: $\pi_S(x, x, x, x, \dots)$ will be written x^ω .

An ω -semigroup is a semigroup equipped with a suitable infinite product.

Definition (ω -semigroup)

An ω -semigroup S is a pair (S_+, S_ω) , where S_+ is a semigroup, S_ω is a set, and equipped with

- an associative mixed product : $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product $\pi_S : S_+^\omega \longrightarrow S_\omega$ satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

Notation: $\pi_S(x, x, x, x, \dots)$ will be written x^ω .

An ω -semigroup is a semigroup equipped with a suitable infinite product.

Definition (ω -semigroup)

An ω -semigroup S is a pair (S_+, S_ω) , where S_+ is a semigroup, S_ω is a set, and equipped with

- an associative mixed product : $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product $\pi_S : S_+^\omega \longrightarrow S_\omega$ satisfying

$$\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) = \pi_S(s_0, s_1, s_2, \dots),$$

$$s \pi_S(s_0, s_1, s_2, \dots) = \pi_S(s, s_0, s_1, s_2, \dots).$$

Notation: $\pi_S(x, x, x, x, \dots)$ will be written x^ω .

An ω -semigroup is a semigroup equipped with a suitable infinite product.

Definition (ω -semigroup)

An ω -semigroup S is a pair (S_+, S_ω) , where S_+ is a semigroup, S_ω is a set, and equipped with

- an associative mixed product : $S_+ \times S_\omega \longrightarrow S_\omega$
- a surjective infinite product $\pi_S : S_+^\omega \longrightarrow S_\omega$ satisfying


$$\begin{aligned}\pi_S(s_0 s_1 \cdots s_{k_1-1}, s_{k_1} \cdots s_{k_2-1}, \dots) &= \pi_S(s_0, s_1, s_2, \dots), \\ s \pi_S(s_0, s_1, s_2, \dots) &= \pi_S(s, s_0, s_1, s_2, \dots).\end{aligned}$$

Notation: $\pi_S(x, x, x, x, \dots)$ will be written x^ω .


Jérémie Cabessa

A *pointed ω -semigroup* is a pair (S, X) , where

- $S = (S_+, S_\omega)$ is an ω -semigroup,
- $X \subseteq S_\omega$.



Among all finite pointed ω -semigroups recognizing a given ω -rational language L , there exists a minimal one, **the syntactic ω -semigroup** of L , denoted by $Synt(L)$.



Example

Consider the language $K = ((a + b)^*a)^\omega$. Then

$\text{Synt}(K) = (S, X)$, where

■ $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ defined by the relations

$$0 \cdot 0 = 0$$

$$0 \cdot 1 = 0$$

$$1 \cdot 0 = 0$$

$$1 \cdot 1 = 1$$

$$00^\omega = 0^\omega$$

$$10^\omega = 0^\omega$$

$$01^\omega = 1^\omega$$

$$11^\omega = 1^\omega$$

■ $X = \{0^\omega\}$.

Example

Consider the language $K = ((a + b)^*a)^\omega$. Then $\text{Synt}(K) = (S, X)$, where

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ defined by the relations

$$\begin{array}{cccc} 0 \cdot 0 = 0 & 0 \cdot 1 = 0 & 1 \cdot 0 = 0 & 1 \cdot 1 = 1 \\ 00^\omega = 0^\omega & 10^\omega = 0^\omega & 01^\omega = 1^\omega & 11^\omega = 1^\omega \end{array}$$

- $X = \{0^\omega\}$.

Example

Consider the language $L = (a\{b, c\}^* \cup \{b\})^\omega$. Then $\text{Synt}(L) = (T, Y)$, where

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$ is defined by

$$a^2 = a$$

$$ab = a$$

$$ac = a$$

$$ba = a$$

$$b^2 = b$$

$$bc = c$$

$$cb = c$$

$$c^2 = c$$

$$b^\omega = a^\omega$$

$$c^\omega = 0$$

$$aa^\omega = a^\omega$$

$$a(ca)^\omega = a^\omega$$

$$ba^\omega = a^\omega$$

$$b(ca)^\omega = (ca)^\omega$$

$$ca^\omega = (ca)^\omega$$

$$c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}$.

Example

Consider the language $L = (a\{b, c\}^* \cup \{b\})^\omega$. Then $\text{Synt}(L) = (T, Y)$, where

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$ is defined by

$$a^2 = a$$

$$ab = a$$

$$ac = a$$

$$ba = a$$

$$b^2 = b$$

$$bc = c$$

$$cb = c$$

$$c^2 = c$$

$$b^\omega = a^\omega$$

$$c^\omega = 0$$

$$aa^\omega = a^\omega$$

$$a(ca)^\omega = a^\omega$$

$$ba^\omega = a^\omega$$

$$b(ca)^\omega = (ca)^\omega$$

$$ca^\omega = (ca)^\omega$$

$$c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}$.

1 We transpose Wadge games on ω -semigroups...

- 1 We transpose Wadge games on ω -semigroups...
- 2 We define the corresponding reduction relation.
- 3 We then describe the resulting hierarchy.

- 1 We transpose Wadge games on ω -semigroups...
- 2 We define the corresponding reduction relation.
- 3 We then describe the resulting hierarchy.

Let $S = (S_+, S_\omega)$ and $T = (T_+, T_\omega)$ be two ω -semigroups, and let also $X \subseteq S_\omega$ and $Y \subseteq T_\omega$.

The infinite two-player game $\text{SG}(X, Y)$ is defined as follows:

- Player I plays elements from S_+ ,
- Player II plays elements from T_+ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let $S = (S_+, S_\omega)$ and $T = (T_+, T_\omega)$ be two ω -semigroups, and let also $X \subseteq S_\omega$ and $Y \subseteq T_\omega$.

The infinite two-player game $\text{SG}(X, Y)$ is defined as follows:

- Player I plays elements from S_+ ,
- Player II plays elements from T_+ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let $S = (S_+, S_\omega)$ and $T = (T_+, T_\omega)$ be two ω -semigroups, and let also $X \subseteq S_\omega$ and $Y \subseteq T_\omega$.

The infinite two-player game $\text{SG}(X, Y)$ is defined as follows:

- Player I plays elements from S_+ ,
- Player II plays elements from T_+ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let $S = (S_+, S_\omega)$ and $T = (T_+, T_\omega)$ be two ω -semigroups, and let also $X \subseteq S_\omega$ and $Y \subseteq T_\omega$.

The infinite two-player game $\text{SG}(X, Y)$ is defined as follows:

- Player I plays elements from S_+ ,
- Player II plays elements from T_+ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,

Let $S = (S_+, S_\omega)$ and $T = (T_+, T_\omega)$ be two ω -semigroups, and let also $X \subseteq S_\omega$ and $Y \subseteq T_\omega$.

The infinite two-player game $\text{SG}(X, Y)$ is defined as follows:

- Player I plays elements from S_+ ,
- Player II plays elements from T_+ ,
- players I and II play alternately, Player I begins,
- Player II may skip her turn, Player I is not allowed,



■ Player II wins the game $\text{SG}(X, Y)$ iff

$$\pi_S(s_0, s_1, \dots) \in X \Leftrightarrow \pi_T(t_0, t_1, \dots) \in Y.$$

Definition (SG-reduction)

We write $X \leq_{\text{SG}} Y$ iff Player II has a winning strategy in $\text{SG}(X, Y)$. And as usual $X \equiv_{\text{SG}} Y$ iff $X \leq_{\text{SG}} Y \leq_{\text{SG}} X$.



■ Player II wins the game $\text{SG}(X, Y)$ iff

$$\pi_S(s_0, s_1, \dots) \in X \Leftrightarrow \pi_T(t_0, t_1, \dots) \in Y.$$

Definition (SG-reduction)

We write $X \leq_{\text{SG}} Y$ iff Player II has a winning strategy in $\text{SG}(X, Y)$. And as usual $X \equiv_{\text{SG}} Y$ iff $X \leq_{\text{SG}} Y \leq_{\text{SG}} X$.



■ Player II wins the game $\text{SG}(X, Y)$ iff

$$\pi_S(s_0, s_1, \dots) \in X \Leftrightarrow \pi_T(t_0, t_1, \dots) \in Y.$$

Definition (SG-reduction)

We write $X \leq_{\text{SG}} Y$ iff Player II has a winning strategy in $\text{SG}(X, Y)$. And as usual $X \equiv_{\text{SG}} Y$ iff $X \leq_{\text{SG}} Y \leq_{\text{SG}} X$.



■ Player II wins the game $\text{SG}(X, Y)$ iff

$$\pi_S(s_0, s_1, \dots) \in X \Leftrightarrow \pi_T(t_0, t_1, \dots) \in Y.$$

Definition (SG-reduction)

We write $X \leq_{\text{SG}} Y$ iff Player II has a winning strategy in $\text{SG}(X, Y)$. And as usual $X \equiv_{\text{SG}} Y$ iff $X \leq_{\text{SG}} Y \leq_{\text{SG}} X$.

Example

Let $S = (\{0, 1\}, \{0 \rightarrow 0, 0 \rightarrow 1, 1 \rightarrow 1\})$ be the ω -semigroup defined by the operations:

$$0 \cdot 0 = 0$$

$$0 \cdot 1 = 0$$

$$1 \cdot 0 = 0$$

$$1 \cdot 1 = 1$$

$$00^\omega = 0 \rightarrow 0$$

$$10^\omega = 0 \rightarrow 0$$

$$01^\omega = 0 \rightarrow 1$$

$$11^\omega = 1 \rightarrow 1$$

- $\pi_S(\alpha) = 0 \rightarrow 0$ iff α contains infinitely many 0's,
- $\pi_S(\alpha) = 0 \rightarrow 1$ iff α contains finitely many 0's,
- $\pi_S(\alpha) = 1 \rightarrow 1$ iff $\alpha = 1^\omega$.

Then $\{1 \rightarrow 1\} \leq_{SG} \{0 \rightarrow 0\}$.

Example

Let $S = (\{0, 1\}, \{0 \rightarrow 0, 0 \rightarrow 1, 1 \rightarrow 1\})$ be the ω -semigroup defined by the operations:

$$0 \cdot 0 = 0$$

$$0 \cdot 1 = 0$$

$$1 \cdot 0 = 0$$

$$1 \cdot 1 = 1$$

$$00^\omega = 0 \rightarrow 0$$

$$10^\omega = 0 \rightarrow 0$$

$$01^\omega = 0 \rightarrow 1$$

$$11^\omega = 1 \rightarrow 1$$

- $\pi_S(\alpha) = 0 \rightarrow 0$ iff α contains infinitely many 0's,
- $\pi_S(\alpha) = 0 \rightarrow 1$ iff α contains finitely many 0's,
- $\pi_S(\alpha) = 1 \rightarrow 1$ iff $\alpha = 1^\omega$.

Then $\{1 \rightarrow 1\} \leq_{SG} \{0 \rightarrow 0\}$.

Example

Let $S = (\{0, 1\}, \{0 \rightarrow 0, 0 \rightarrow 1, 1 \rightarrow 1\})$ be the ω -semigroup defined by the operations:

$$0 \cdot 0 = 0$$

$$0 \cdot 1 = 0$$

$$1 \cdot 0 = 0$$

$$1 \cdot 1 = 1$$

$$00^\omega = 0 \rightarrow 0$$

$$10^\omega = 0 \rightarrow 0$$

$$01^\omega = 0 \rightarrow 1$$

$$11^\omega = 1 \rightarrow 1$$

- $\pi_S(\alpha) = 0 \rightarrow 0$ iff α contains infinitely many 0's,
- $\pi_S(\alpha) = 0 \rightarrow 1$ iff α contains finitely many 0's,
- $\pi_S(\alpha) = 1 \rightarrow 1$ iff $\alpha = 1^\omega$.

Then $\{1 \rightarrow 1\} \leq_{SG} \{0 \rightarrow 0\}$.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 1 $1 \rightarrow 1 \in \{1 \rightarrow 1\}$

Player II 0 0 0 $0 \rightarrow 0 \in \{0 \rightarrow 0\}$

Therefore Player II wins.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Example (continued)

We give a w. s. for Player II in $\text{SG}(\{1 \rightarrow 1\}, \{0 \rightarrow 0\})$:

Player I 1 1 0 $0 \rightarrow 0$ or $0 \rightarrow 1 \notin \{1 \rightarrow 1\}$

Player II 0 0 1's until the end $0 \rightarrow 1 \notin \{0 \rightarrow 0\}$

Therefore Player II wins again.

Definition (FSG-hierarchy)

The collection of finite pointed ω -semigroups ordered by the \leq_{SG} -relation is called *the FSG-hierarchy*.

Proposition (SG-Borel Determinacy)

Let (S, X) and (T, Y) be two finite pointed ω -semigroups. Then the game $SG(X, Y)$ is determined.

Proof.

A consequence of Borel Wadge determinacy. A given player has a w. s. in $SG(X, Y)$ iff this same player has a w. s. in $W(\pi_S^{-1}(X), \pi_T^{-1}(Y))$. □

Definition (FSG-hierarchy)

The collection of finite pointed ω -semigroups ordered by the \leq_{SG} -relation is called *the FSG-hierarchy*.

Proposition (SG-Borel Determinacy)

Let (S, X) and (T, Y) be two finite pointed ω -semigroups. Then the game $SG(X, Y)$ is determined.

Proof.

A consequence of Borel Wadge determinacy. A given player has a w. s. in $SG(X, Y)$ iff this same player has a w. s. in $W(\pi_S^{-1}(X), \pi_T^{-1}(Y))$. □

Definition (FSG-hierarchy)

The collection of finite pointed ω -semigroups ordered by the \leq_{SG} -relation is called *the FSG-hierarchy*.

Proposition (SG-Borel Determinacy)

Let (S, X) and (T, Y) be two finite pointed ω -semigroups. Then the game $SG(X, Y)$ is determined.

Proof.

A consequence of Borel Wadge determinacy. A given player has a w. s. in $SG(X, Y)$ iff this same player has a w. s. in $W(\pi_S^{-1}(X), \pi_T^{-1}(Y))$.



Corollary

- *The \leq_{SG} -antichains have length at most two.*
- *If X and Y are incomparable, then $X \equiv_{SG} Y^c$.*
- *The partial ordering \leq_{SG} is wellfounded on finite pointed ω -semigroups.*


Corollary

- *The \leq_{SG} -antichains have length at most two.*
- *If X and Y are incomparable, then $X \equiv_{SG} Y^c$.*
- *The partial ordering \leq_{SG} is wellfounded on finite pointed ω -semigroups.*

Corollary


- *The \leq_{SG} -antichains have length at most two.*
- *If X and Y are incomparable, then $X \equiv_{SG} Y^c$.*
- *The partial ordering \leq_{SG} is wellfounded on finite pointed ω -semigroups.*

The FSG-hierarchy



Theorem

The FSG-hierarchy and the Wagner hierarchy are isomorphic.




Corollary

The FSG-hierarchy has height ω^ω , and it is decidable.

Theorem


The FSG-hierarchy and the Wagner hierarchy are isomorphic.



Corollary

The FSG-hierarchy has height ω^ω , and it is decidable.

Completely algebraic decidability procedure: **one can compute the Wagner degree of an ω -rational language directly on its syntactic image.**



Example

Consider the syntactic pointed ω -semigroup of $K = ((a + b)^*a)^\omega$:

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$ defined by the relations

$$\begin{array}{cccc} 0 \cdot 0 = 0 & 0 \cdot 1 = 0 & 1 \cdot 0 = 0 & 1 \cdot 1 = 1 \\ 00^\omega = 0^\omega & 10^\omega = 0^\omega & 01^\omega = 1^\omega & 11^\omega = 1^\omega \end{array}$$

- $X = \{0^\omega\}$.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$

$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$

●

1

●

0

Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

■ $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$


$0 \cdot 0 = 0$ $0 \cdot 1 = 0$

$1 \cdot 0 = 0$ $1 \cdot 1 = 1$

$00^\omega = 0^\omega$ $10^\omega = 0^\omega$

$01^\omega = 1^\omega$ $11^\omega = 1^\omega$

■ $X = \{0^\omega\}$.



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$


$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

- $S = (\{0, 1\}, \{0^\omega, 1^\omega\})$
 $0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$
 $1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$
 $00^\omega = 0^\omega \quad 10^\omega = 0^\omega$
 $01^\omega = 1^\omega \quad 11^\omega = 1^\omega$
- $X = \{0^\omega\}$.



Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example (continued)

$$\blacksquare S = (\{0, 1\}, \{0^\omega, 1^\omega\})$$


$$0 \cdot 0 = 0 \quad 0 \cdot 1 = 0$$

$$1 \cdot 0 = 0 \quad 1 \cdot 1 = 1$$

$$00^\omega = 0^\omega \quad 10^\omega = 0^\omega$$

$$01^\omega = 1^\omega \quad 11^\omega = 1^\omega$$

$$\blacksquare X = \{0^\omega\}.$$



Therefore $d_{SG}((S, X)) = d_W(K) = \omega$, and K is non-self-dual.

Example

Consider the syntactic pointed ω -semigroup of

$$L = (a\{b, c\}^* \cup \{b\})^\omega:$$

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$ is defined by

$$a^2 = a$$

$$ab = a$$

$$ac = a$$

$$ba = a$$

$$b^2 = b$$

$$bc = c$$

$$cb = c$$

$$c^2 = c$$

$$b^\omega = a^\omega$$

$$c^\omega = 0$$

$$aa^\omega = a^\omega$$

$$a(ca)^\omega = a^\omega$$

$$ba^\omega = a^\omega$$

$$b(ca)^\omega = (ca)^\omega$$

$$ca^\omega = (ca)^\omega$$

$$c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$

Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$


$$b^2 = b$$

$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0 \quad b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$



Example (continued)

■ $T = (\{a, b, c, ca\}, \{a^\omega, (ca)^\omega, 0\})$

$$a^2 = ab = ac = ba = a$$

$$b^2 = b$$


$$bc = cb = c^2 = c$$

$$b^\omega = aa^\omega = a(ca)^\omega = ba^\omega = a^\omega$$

$$c^\omega = 0$$

$$b(ca)^\omega = ca^\omega = c(ca)^\omega = (ca)^\omega$$

■ $Y = \{a^\omega\}.$




Therefore $d_{SG}((T, Y)) = d_W(L) = \omega^2$,
and L is non-self-dual.

Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I

(T,Y) II

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1

(T,Y) II

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1

(T,Y) II a

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1

(T,Y) II a

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1

(T,Y) II a c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1

(T,Y) II a c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1

(T,Y) II a c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1

(T,Y) II a c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1

(T,Y) II a c c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1

(T,Y) II a c c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1

(T,Y) II a c c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0

(T,Y) II a c c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0

(T,Y) II a c c -

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1

(T,Y) II a c c -

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1

(T,Y) II a c c - c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1

(T,Y) II a c c - c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1

(T,Y) II a c c - c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1 0

(T,Y) II a c c - c

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1 0

(T,Y) II a c c - c ca

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1 0

(T,Y) II a c c - c ca

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1 0

(T,Y) II a c c - c ca

Player I



Player II




Therefore $(S, X) \leq_{SG} (T, Y)$, or equivalently $K \leq_W L$.


(S,X) I 1 1 1 0 1 0

(T,Y) II a c c - c ca

Player I



Player II



Summary:

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- One can compute the Wagner degree of an ω -rational language directly on its syntactic image.
- Build a finite pointed ω -semigroup of any given Wagner degree.

Summary:

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- One can compute the Wagner degree of an ω -rational language directly on its syntactic image.
- Build a finite pointed ω -semigroup of any given Wagner degree.

Summary:

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- **One can compute the Wagner degree of an ω -rational language directly on its syntactic image.**
- Build a finite pointed ω -semigroup of any given Wagner degree.

Summary:

- The FSG-hierarchy is the algebraic counterpart of the Wagner hierarchy.
- Decidability procedure of the FSG-hierarchy.
- **One can compute the Wagner degree of an ω -rational language directly on its syntactic image.**
- Build a finite pointed ω -semigroup of any given Wagner degree.

Future work:

- Computer science: towards a simmlar analysis of the algebraic counterpart of more complex abstract machines.
- Descriptive set theory: towards an algebraic study of the Wadge hierarchy.

Future work:

- Computer science: towards a simmlar analysis of the algebraic counterpart of more complex abstract machines.
- Descriptive set theory: towards an algebraic study of the Wadge hierarchy.

Future work:

- Computer science: towards a simmlar analysis of the algebraic counterpart of more complex abstract machines.
- Descriptive set theory: towards an algebraic study of the Wadge hierarchy.